

*This copy is for your personal, non-commercial use only.*

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines [here](#).

**The following resources related to this article are available online at [www.sciencemag.org](http://www.sciencemag.org) (this information is current as of May 28, 2010):**

**Updated information and services**, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/cgi/content/full/326/5958/1397>

**Supporting Online Material** can be found at:

<http://www.sciencemag.org/cgi/content/full/1180251/DC1>

This article **cites 14 articles**, 5 of which can be accessed for free:

<http://www.sciencemag.org/cgi/content/full/326/5958/1397#otherarticles>

This article has been **cited by** 3 article(s) on the ISI Web of Science.

This article has been **cited by** 1 articles hosted by HighWire Press; see:

<http://www.sciencemag.org/cgi/content/full/326/5958/1397#otherarticles>

This article appears in the following **subject collections**:

Ecology

<http://www.sciencemag.org/cgi/collection/ecology>

Science and Policy

[http://www.sciencemag.org/cgi/collection/sci\\_policy](http://www.sciencemag.org/cgi/collection/sci_policy)

- G. Johnson, W. Ruddiman, Eds. (Ocean Drilling Program, College Station, TX, 1996), pp. 515–568.
25. K. Billups, D. Schrag, *Earth Planet. Sci. Lett.* **209**, 181 (2003).
26. A. Krylov *et al.*, *Paleoceanography* **23**, PA1506 (2008).
27. A. R. Lewis *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **105**, 10676 (2008).
28. T. Naish *et al.*, *Nature* **458**, 322 (2009).
29. B. Flower, J. Kennett, *Paleoceanography* **10**, 1095 (1995).
30. T. Naish *et al.*, paper presented at the International Symposium on Antarctic Earth Sciences, Santa Barbara, CA, 26 to 31 August 2007.
31. D. J. Lunt, G. L. Foster, A. M. Haywood, E. J. Stone, *Nature* **454**, 1102 (2008).
32. L. E. Lisiecki, M. E. Raymo, *Paleoceanography* **20**, PA1003 (2005).
33. L. Krissiek, in *Proceedings of the Ocean Drilling Program, Scientific Results*, vol. 145, D. Rea, I. Basov, D. Scholl, J. Allan, Eds. (Ocean Drilling Program, College Station, TX, 1995), pp. 179–194.
34. E. Cowan, in *Proceedings of the Ocean Drilling Program, Scientific Results*, vol. 178, P. Barker, A. Camerlenghi, G. Acton, Eds. (Ocean Drilling Program, College Station, TX, 2001), pp. 1–22.
35. R. M. DeConto *et al.*, *Nature* **455**, 652 (2008).
36. We thank K. Caldeira, H. Elderfield, J. Eiler, T. Naish, D. Sigman, anonymous reviewers, and the editor for their comments on this work, which substantially improved the manuscript. We also thank J. Booth, E. Khadun, O. Shorttle, L. Thanalasundaram, and A. Bufe for invaluable assistance with sample preparation; L. Booth, J. Day, and M. Greaves (supported by grant NE/F004966/1) for technical assistance; L. Lisiecki for assistance with the age model; and S. Crowhurst, A. Gagnon, S. John, N. Meckler, B. Passey, N. Thiagarajan, and J. Yu for discussing this work. Support was provided to A.K.T. by UCLA, National Environmental Research Council (NERC) (fellowship NE/D009049/1), and Magdalene College; to C.D.R. by NERC (studentship NER/S/A/2006/14070); and to R.A.E. by a Caltech Chancellors Postdoctoral Scholarship. Samples for this study were obtained from the Godwin Laboratory sample archives and the Ocean Drilling Program.

### Supporting Online Material

www.sciencemag.org/cgi/content/full/1178296/DC1  
Materials and Methods

Figs. S1 to S9

Tables S1 to S12

References

26 June 2009; accepted 28 September 2009

Published online 8 October 2009;

10.1126/science.1178296

Include this information when citing this paper.

## Indirect Emissions from Biofuels: How Important?

Jerry M. Melillo,<sup>1\*</sup> John M. Reilly,<sup>2</sup> David W. Kicklighter,<sup>1</sup> Angelo C. Gurgel,<sup>2,3</sup> Timothy W. Cronin,<sup>1,2</sup> Sergey Paltsev,<sup>2</sup> Benjamin S. Felzer,<sup>1,4</sup> Xiaodong Wang,<sup>2,5</sup> Andrei P. Sokolov,<sup>2</sup> C. Adam Schlosser<sup>2</sup>

A global biofuels program will lead to intense pressures on land supply and can increase greenhouse gas emissions from land-use changes. Using linked economic and terrestrial biogeochemistry models, we examined direct and indirect effects of possible land-use changes from an expanded global cellulosic bioenergy program on greenhouse gas emissions over the 21st century. Our model predicts that indirect land use will be responsible for substantially more carbon loss (up to twice as much) than direct land use; however, because of predicted increases in fertilizer use, nitrous oxide emissions will be more important than carbon losses themselves in terms of warming potential. A global greenhouse gas emissions policy that protects forests and encourages best practices for nitrogen fertilizer use can dramatically reduce emissions associated with biofuels production.

Expanded use of bioenergy causes land-use changes and increases in terrestrial carbon emissions (1, 2). The recognition of this has led to efforts to determine the credit toward meeting low carbon fuel standards (LCFS) for different forms of bioenergy with an accounting of direct land-use emissions as well as emissions from land use indirectly related to bioenergy production (3, 4). Indirect emissions occur when biofuels production on agricultural land displaces agricultural production and causes additional land-use change that leads to an increase in net greenhouse gas (GHG) emissions (2, 4). The control of GHGs through a cap-and-trade or tax policy, if extended to include emissions (or credits for uptake) from land-use change combined with monitoring of carbon stored in vegetation and soils and enforcement of such

policies, would eliminate the need for such life-cycle accounting (5, 6). There are a variety of concerns (5) about the practicality of including land-use change emissions in a system designed to reduce emissions from fossil fuels, and that may explain why there are no concrete proposals in major countries to do so. In this situation, fossil energy control programs (LCFS or carbon taxes) must determine how to treat the direct and indirect GHG emissions associated with the carbon intensity of biofuels.

The methods to estimate indirect emissions remain controversial. Quantitative analyses to date have ignored these emissions (1), considered those associated with crop displacement from a limited area (2), confounded these emissions with direct or general land-use emissions (6–8), or developed estimates in a static framework of today's economy (3). Missing in these analyses is how to address the full dynamic accounting of biofuel carbon intensity (CI), which is defined for energy as the GHG emissions per megajoule of energy produced (9), that is, the simultaneous consideration of the potential of net carbon uptake through enhanced management of poor or degraded lands, nitrous oxide (N<sub>2</sub>O) emissions that would accompany increased use of fertilizer, environmental effects on terrestrial carbon storage [such as climate

change, enhanced carbon dioxide (CO<sub>2</sub>) concentrations, and ozone pollution], and consideration of the economics of land conversion. The estimation of emissions related to global land-use change, both those on land devoted to biofuel crops (direct emissions) and those indirect changes driven by increased demand for land for biofuel crops (indirect emissions), requires an approach to attribute effects to separate land uses.

We applied an existing global modeling system that integrates land-use change as driven by multiple demands for land and that includes dynamic greenhouse gas accounting (10, 11). Our modeling system, which consists of a computable general equilibrium (CGE) model of the world economy (10, 12) combined with a process-based terrestrial biogeochemistry model (13, 14), was used to generate global land-use scenarios and explore some of the environmental consequences of an expanded global cellulosic biofuels program over the 21st century. The biofuels scenarios we focus on are linked to a global climate policy to control GHG emissions from industrial and fossil fuel sources that would, absent feedbacks from land-use change, stabilize the atmosphere's CO<sub>2</sub> concentration at 550 parts per million by volume (ppmv) (15). The climate policy makes the use of fossil fuels more expensive, speeds up the introduction of biofuels, and ultimately increases the size of the biofuel industry, with additional effects on land use, land prices, and food and forestry production and prices (16).

We considered two cases in order to explore future land-use scenarios: Case 1 allows the conversion of natural areas to meet increased demand for land, as long as the conversion is profitable; case 2 is driven by more intense use of existing managed land. To identify the total effects of biofuels, each of the above cases is compared with a scenario in which expanded biofuel use does not occur (16). In the scenarios with increased biofuels production, the direct effects (such as changes in carbon storage and N<sub>2</sub>O emissions) are estimated only in areas devoted to biofuels. Indirect effects are defined as the differences between the total effects and the direct effects.


At the beginning of the 21st century, ~31.5% of the total land area (133 million km<sup>2</sup>) was in

<sup>1</sup>The Ecosystems Center, Marine Biological Laboratory (MBL), 7 MBL Street, Woods Hole, MA 02543, USA. <sup>2</sup>Joint Program on the Science and Policy of Global Change, Massachusetts Institute of Technology (MIT), 77 Massachusetts Avenue, MIT E19-411, Cambridge, MA 02139-4307, USA. <sup>3</sup>Department of Economics, University of São Paulo, Ribeirão Preto 4EES, Brazil.

<sup>4</sup>Department of Earth and Environmental Sciences, Lehigh University, 31 Williams Drive, Bethlehem, PA 18015, USA.


<sup>5</sup>School of Public Administration, Zhejiang University, Hangzhou 310000, Zhejiang Province, People's Republic of China (PRC).

\*To whom correspondence should be addressed. E-mail: jmelillo@mbll.edu




**Fig. 1.** Projected changes in global land cover for land-use case 1 (A) and case 2 (B). In either case, biofuels supply most of the world’s liquid fuel needs by 2100. In case 1, 365 EJ of biofuel is produced in 2100, using 16.2% (21.6 million km<sup>2</sup>) of the total land area; natural forest area declines from 34.4 to 15.1 million km<sup>2</sup> (56%), and pasture area declines from 25.8 to 22.1 million km<sup>2</sup> (14%). In case 2, 323 EJ of biofuels are produced in 2100, using 20.6 million km<sup>2</sup> of land; pasture areas decrease by 10.3 million km<sup>2</sup> (40%), and forest area declines by 8.4 million km<sup>2</sup> (24% of forest area). Simulations show that these major land-use changes will take place in the tropics and subtropics, especially in Africa and the Americas (fig. S2).

**Fig. 2.** Partitioning of direct (dark gray) and indirect effects (light gray) on projected cumulative land carbon flux since the year 2000 (black line) from cellulosic biofuel production for land-use case 1 (A) and case 2 (B). Positive values represent carbon sequestration, whereas negative values represent carbon emissions by land ecosystems. In case 1, the cumulative loss is 92 Pg CO<sub>2</sub>eq by 2100, with the maximum loss (164 Pg CO<sub>2</sub>eq) occurring in the 2050 to 2055 time frame, indirect losses of 110 Pg CO<sub>2</sub>eq, and direct losses of 54 Pg CO<sub>2</sub>eq. In the second half of the century, there is net accumulation of 72 Pg CO<sub>2</sub>eq mostly in the soil in response to the use of nitrogen fertilizers. In case 2, land areas are projected to have a net accumulation of 75 Pg CO<sub>2</sub>eq as a result of biofuel production, with maximum loss of 26 Pg CO<sub>2</sub>eq in the 2035 to 2040 time frame, followed by substantial accumulation.



**Fig. 3.** Partitioning of greenhouse gas balance since the year 2000 (black line) as influenced by cellulosic biofuel production for land-use case 1 (A) and case 2 (B) among fossil fuel abatement (yellow), net land carbon flux (blue), and fertilizer N<sub>2</sub>O emissions (red). Positive values are abatement benefits, and negative values are emissions. Net land carbon flux is the same as in Fig. 2. For case 1, N<sub>2</sub>O emissions over the century are 286 Pg CO<sub>2</sub>eq; for case 2, N<sub>2</sub>O emissions are 238 Pg CO<sub>2</sub>eq.



**Table 1.** Carbon intensity index associated with cellulosic biofuel production for two land-use scenario cases. Units are g CO<sub>2</sub>eq MJ<sup>-1</sup>, with negative values indicating carbon accumulation.

Variable	Case 1			Case 2		
Time period	2000–2030	2000–2050	2000–2100	2000–2030	2000–2050	2000–2100
Direct land C	11	27	0	-52	-24	-7
Indirect land C	190	57	7	181	31	1
Fertilizer N <sub>2</sub> O	29	28	20	30	26	19
Total	229	112	26	158	32	13

agriculture: 12.1% (16.1 million km<sup>2</sup>) in crops and 19.4% (25.8 million km<sup>2</sup>) in pasture (17). In both cases of increased biofuels use, land devoted to biofuels becomes greater than all area currently devoted to crops by the end of the 21st century, but in case 2 less forest land is converted (Fig. 1). Changes in net land fluxes are also associated with how land is allocated for biofuels production (Fig. 2). In case 1, there is a larger loss of carbon than in case 2, especially at mid-century. Indirect land use is responsible for substantially greater carbon losses than direct land use in both cases during the first half of the century. In both cases, there is carbon accumulation in the latter part of the century. The estimates include CO<sub>2</sub> from burning and decay of vegetation and slower release of carbon as CO<sub>2</sub> from disturbed soils. The estimates also take into account reduced carbon sequestration capacity of the cleared areas, including that which would have been stimulated by increased ambient CO<sub>2</sub> levels. Smaller losses in the early years in case 2 are due to less deforestation and more use of pasture, shrubland, and savanna, which have lower carbon stocks than forests and, once under more intensive management, accumulate soil carbon. Much of the soil carbon accumulation is projected to occur in sub-Saharan Africa, an attractive area for growing biofuels in our economic analyses because the land is relatively inexpensive (10) and simple management interventions such as fertilizer additions can dramatically increase crop productivity (18).

Estimates of land devoted to biofuels in our two scenarios (15 to 16%) are well below the estimate of ~50% in a recent analysis (6) that does not control land-use emissions. The higher number is based on an analysis that has a lower concentration target (450 ppmv CO<sub>2</sub>), does not account for price-induced intensification of land use, and does not explicitly consider concurrent changes in other environmental factors. In analyses that include land-use emissions as part of the policy (6–8), less area is estimated to be devoted to biofuels (3 to 8%).

The carbon losses associated with the combined direct and indirect biofuel emissions estimated for our case 1 are similar to a previous estimate (7), which shows larger losses of carbon per unit area converted to biofuels production. These larger losses per unit area result from a combination of factors, including a greater simulated response of plant productivity to changes in climate and atmospheric CO<sub>2</sub> (15) and the lack of any negative effects on plant productivity of elevated tropospheric ozone (19, 20).

We also simulated the emissions of N<sub>2</sub>O from additional fertilizer that would be required to grow biofuel crops. Over the century, the N<sub>2</sub>O emissions become larger in CO<sub>2</sub> equivalent (CO<sub>2</sub>eq) than carbon emissions from land use (Fig. 3). The net GHG effect of biofuels also changes over time; for case 1, the net GHG balance is -90 Pg CO<sub>2</sub>eq through 2050 (a negative sign indicates a source; a positive sign indicates a sink), whereas it is +579 through

2100. For case 2, the net GHG balance is +57 Pg CO<sub>2</sub>eq through 2050 and +679 through 2100. We estimate that by the year 2100, biofuels production accounts for about 60% of the total annual N<sub>2</sub>O emissions from fertilizer application in both cases, where the total for case 1 is 18.6 Tg N yr<sup>-1</sup> and for case 2 is 16.1 Tg N yr<sup>-1</sup>. These total annual land-use N<sub>2</sub>O emissions are about 2.5 to 3.5 times higher than comparable estimates from an earlier study (8). Our larger estimates result from differences in the assumed proportion of nitrogen fertilizer lost as N<sub>2</sub>O (21) as well as differences in the amount of land devoted to food and biofuel production. Best practices for the use of nitrogen fertilizer, such as synchronizing fertilizer application with plant demand (22), can reduce N<sub>2</sub>O emissions associated with biofuels production.

The CI of fuel was also calculated across three time periods (Table 1) so as to compare with displaced fossil energy in a LCFS and to identify the GHG allowances that would be required for biofuels in a cap-and-trade program. Previous CI estimates for California gasoline (3) suggest that values less than ~96 g CO<sub>2</sub>eq MJ<sup>-1</sup> indicate that blending cellulosic biofuels will help lower the carbon intensity of California fuel and therefore contribute to achieving the LCFS. Entries that are higher than 96 g CO<sub>2</sub>eq MJ<sup>-1</sup> would raise the average California fuel carbon intensity and thus be at odds with the LCFS. Therefore, the CI values for case 1 are only favorable for biofuels if the integration period extends into the second half of the century. For case 2, the CI values turn favorable for biofuels over an integration period somewhere between 2030 and 2050. In both cases, the CO<sub>2</sub> flux has approached zero by the end of the century when little or no further land conversion is occurring and emissions from decomposition are approximately balancing carbon added to the soil from unharvested components of the vegetation (roots). Although the carbon accounting ends up as a nearly net neutral effect, N<sub>2</sub>O emissions continue. Annual estimates start high, are variable from year to year because they depend on climate, and generally decline over time.

One of the perplexing issues for policy analysts has been predicting the dynamics of the CI over different integration periods [supporting online material (SOM) text]. If one integrates over a long enough period, biofuels show a substantial greenhouse gas advantage, but over a short period they have a higher CI than fossil fuel (3). Drawing on previous analyses (5, 23), we argue that a solution need not be complex and can avoid valuing climate damages by using the immediate (annual) emissions (direct and indirect) for the CI calculation. In other words, CI estimates should not integrate over multiple years but rather simply consider the fuel offset for the policy time period (normally a single year). This becomes evident in case 1. Despite the promise of eventual long-term economic benefits, a substantial penalty—in fact, possibly worse than with gasoline—in the first few decades may render

the near-term cost of the carbon debt difficult to overcome in this case.

In case 2, in which there is less willingness to convert land, the economics of biofuels would be favorable sooner. Greater measures to protect forests could make the economics and CI of biofuels even more favorable because improved management on low-quality or degraded land can lead to carbon accumulation in the soil rather than a carbon loss (fig. S3). Our results suggest that tropical regions that are currently suffering substantial amounts of deforestation may also be the most competitive producers of biofuels. Our suggested strategy of not integrating over future fuel offsets increases the near-term CI of biofuels unless forested lands globally are better protected. Success in avoiding deforestation will be reflected in lower estimates of indirect emissions and lower carbon penalties.

#### References and Notes

1. J. Fargione *et al.*, *Science* **319**, 1235 (2008).
2. T. Searchinger *et al.*, *Science* **319**, 1238 (2008).
3. California Environmental Protection Agency (EPA), "Proposed regulation to implement the low carbon fuel standard," vol. 1 (EPA, Sacramento, CA, 2009); [www.arb.ca.gov/fuels/lcfs/030409lcfs\\_isor\\_vol1.pdf](http://www.arb.ca.gov/fuels/lcfs/030409lcfs_isor_vol1.pdf).
4. E. Gallagher, *The Gallagher Review of the Indirect Effects of Biofuels Production* (Renewable Fuels Agency, East Sussex, UK, 2008).
5. J. Reilly *et al.*, in *Greenhouse Gas Sinks*, D. S. Reay, C. N. Hewitt, K. A. Smith, J. Grace, Eds. (CABI Publishing, Wallingford, UK, 2007), chap 8, pp. 115–142.
6. M. Wise *et al.*, *Science* **324**, 1183 (2009).
7. R. Leemans, A. van Amstel, C. Battjes, E. Kreileman, S. Toet, *Glob. Environ. Change* **6**, 335 (1996).
8. B. Strengers, R. Leemans, B. Eickhout, B. de Vries, L. Bouwman, *GeoJournal* **61**, 381 (2004).
9. M. R. Raupach *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **104**, 10288 (2007).

10. A. Gurgel, J. M. Reilly, S. Paltsev, *J. Agric. Food Ind. Org.* [www.bepress.com/jafio/vol5/iss2/art9](http://www.bepress.com/jafio/vol5/iss2/art9).
11. J. Melillo *et al.*, "Unintended environmental consequences of a global biofuels program" [Report 168, MIT Joint Program on the Science and Policy of Global Change (JPSPGC), Cambridge, 2009]; [http://globalchange.mit.edu/files/document/MITJPSPGC\\_Rpt168.pdf](http://globalchange.mit.edu/files/document/MITJPSPGC_Rpt168.pdf).
12. S. Paltsev *et al.*, "The MIT emissions prediction and policy analysis (EPPA) model: version 4" (MIT JPSPGC, Cambridge, 2005); [http://globalchange.mit.edu/files/document/MITJPSPGC\\_Rpt125.pdf](http://globalchange.mit.edu/files/document/MITJPSPGC_Rpt125.pdf).
13. J. Melillo *et al.*, *Nature* **363**, 234 (1993).
14. B. Felzer *et al.*, *Tellus* **56B**, 230 (2004).
15. A. Sokolov *et al.*, *J. Clim.* **21**, 3776 (2008).
16. Materials and methods are available as supporting material on Science Online.
17. G. Hurtt *et al.*, *Glob. Change Biol.* **12**, 1208 (2006).
18. P. A. Sanchez, *Science* **295**, 2019 (2002).
19. B. Felzer *et al.*, *Clim. Change* **73**, 345 (2005).
20. X. Wang, thesis, MIT (2008); available at [http://globalchange.mit.edu/files/document/Wang\\_PhD\\_08.pdf](http://globalchange.mit.edu/files/document/Wang_PhD_08.pdf).
21. P. J. Crutzen, A. R. Mosier, K. A. Smith, W. Winiwarter, *Atmos. Chem. Phys.* **8**, 389 (2008).
22. G. P. Robertson, in *Ecology in Agriculture*, L. Jackson, Ed. (Academic Press, New York, 1997), pp. 347–365.
23. H. Herzog *et al.*, *Clim. Change* **59**, 293 (2003).
24. This research was supported in part by the David and Lucile Packard Foundation to the MBL, Department of Energy, Office of Science (BER) grants DE-FG02-94ER61937, DE-FG02-93ER61677, and DE-FG02-08ER64648; EPA grant XA-83240101; NSF grant BCS-0410344; and the industrial and foundation sponsors of the MIT JPSPGC.

#### Supporting Online Material

[www.sciencemag.org/cgi/content/full/1180251/DC1](http://www.sciencemag.org/cgi/content/full/1180251/DC1)  
Materials and Methods  
SOM Text  
Figs. S1 to S3  
Tables S1 to S6  
References

6 August 2009; accepted 7 October 2009  
Published online 22 October 2009;  
10.1126/science.1180251  
Include this information when citing this paper.

## Elevated CO<sub>2</sub> Reduces Losses of Plant Diversity Caused by Nitrogen Deposition

Peter B. Reich

The interactive effects of rising atmospheric carbon dioxide (CO<sub>2</sub>) concentrations and elevated nitrogen (N) deposition on plant diversity are not well understood. This is of concern because both factors are important components of global environmental change and because each might suppress diversity, with their combined effects possibly additive or synergistic. In a long-term open-air experiment, grassland assemblages planted with 16 species were grown under all combinations of ambient and elevated CO<sub>2</sub> and ambient and elevated N. Over 10 years, elevated N reduced species richness by 16% at ambient CO<sub>2</sub> but by just 8% at elevated CO<sub>2</sub>. This resulted from multiple effects of CO<sub>2</sub> and N on plant traits and soil resources that altered competitive interactions among species. Elevated CO<sub>2</sub> thus ameliorated the negative effects of N enrichment on species richness.

Two global change factors likely to have widespread influence on plant communities are nitrogen (N) deposition and rising atmospheric carbon dioxide (CO<sub>2</sub>) levels (1–7). Levels of N deposition and CO<sub>2</sub> have risen in recent decades and are expected to increase further (8). Because increased CO<sub>2</sub> and N supply

often drive plant stoichiometry in opposite directions but productivity in the same direction, and as plant resources are primarily available above-

Department of Forest Resources, University of Minnesota, 1530 Cleveland Avenue North, St. Paul, MN 55108, USA. E-mail: preich@umn.edu