

1 **POTENTIAL IMPACTS OF CLIMATE CHANGE TO FISHER HABITAT IN**  
2 **CALIFORNIA: A PRELIMINARY ASSESSMENT**

3  
4 Hugh D. Safford  
5 USDA-Forest Service  
6 Pacific Southwest Region  
7 1323 Club Drive  
8 Vallejo, CA 94592  
9 hughsafford@fs.fed.us

10  
11  
12 **ABSTRACT**

13  
14 An assessment is made of the potential impacts of climate change on the habitat of the California  
15 fisher (*Martes pennanti*). Previous climate change and ecosystem response modeling results are  
16 combined with forest growth and disturbance simulations to provide insight into the likely  
17 outcomes of warming climates on vegetation floristics and structure in forests currently inhabited  
18 by fisher. In California, hardwoods (primarily oaks) have been found to be an extremely  
19 important source of denning and resting structures for fisher. GCM-coupled ecosystem modeling  
20 predicts that climate warming will drive an increase in the hardwood component of montane  
21 forests in California, especially where and if precipitation increases. Fire is predicted to become  
22 more frequent, and in some cases more intense, across much of the fisher's range as well. Greater  
23 activity of fire may negatively impact some of the structural habitat features that are key to fisher  
24 occurrence, such as large tree densities and high canopy cover. Rates of snag and surface woody  
25 debris creation may go up in the short term, but greatly increased frequencies of fire may  
26 ultimately reduce these important structural features over the long term. Little or nothing has  
27 been published on direct metabolic impacts of warming climates on fisher or its prey, and these  
28 types of effects may prove more important to future fisher distribution than indirect effects  
29 mediated through changing habitats. Complicated and synergistic interactions between factors  
30 are likely, and make any precise predictions of climate change impacts on fisher currently  
31 impossible.

32  
33 Key words: California, climate change, fire, fisher, hardwoods, mixed conifer forest

34  
35 **INTRODUCTION**

36  
37 *The fisher and its habitat*

38  
39 Fisher (*Martes pennanti*) is a mesocarnivore belonging to the mink family (Mustelidae). Fishers  
40 are habitat specialists and are considered to be among the western North American carnivore  
41 species most vulnerable to human disturbance and habitat alteration (Powell and Zielinski 1999,  
42 Harrison et al. 2004, Zielinski et al. 2005). Evidence is that fishers once ranged throughout the  
43 lower montane and montane zones of the Klamath Ranges and the Sierra Nevada (Zielinski et al.  
44 2005), but today they are restricted to two disjunct populations separated by more than 400 km  
45 (Figure 1). Home ranges of fishers in California vary from 4-90 km<sup>2</sup>, with males having  
46 significantly larger ranges; averages are about 15 km<sup>2</sup> for females and 40 km<sup>2</sup> for males (Powell  
47 and Zielinski 1994). In California, fishers are found from near sea level in the northwest, to as  
48 high as 2700 m (and above) in the Sierra Nevada; females generally remain at lower elevations  
49 than males (Zielinski et al. 2004a, 2005). The maximum documented dispersal distance for  
50 individual California fishers is about 100 km (Zielinski et al. 2005).

51

1 Floristic and structural characteristics of forests supporting fisher have been described by Powell  
2 (1993), Powell and Zielinski (1994), Seglund (1995), Klug (1997), Carroll et al. (1999), Mazzoni  
3 2002, Zielinski et al. (2004a,b; 2006), Carroll (2005), and Yaeger (2005). In both California  
4 population areas, midseral conifer and mixed conifer-hardwood forests comprise the bulk of  
5 fisher habitat. In the Klamaths, fisher home ranges are primarily in Douglas-fir (*Pseudotsuga*  
6 *menziesii*) and/or white fir (*Abies concolor*) forest, with an important component of conifer-oak  
7 or pure oak stands; in the southern Sierra Nevada, the preferred floristic conditions range from  
8 mixed conifer to ponderosa pine (*Pinus ponderosa*) to montane hardwood. Oaks – chiefly black  
9 oak (*Quercus kelloggii*) - appear to be a key component of the habitat, especially in the southern  
10 Sierra Nevada (Carroll et al. 1999, Zielinski et al. 2004a). Forest structural characteristics within  
11 fisher home ranges are strongly skewed toward mid- to late-seral stands with high canopy cover;  
12 large, cavity-forming trees are required for resting and denning habitat (Seglund 1995, Zielinski  
13 et al. 2004b, Yaeger 2005). Geographic conditions correlated with core fisher habitat in  
14 California include complex topography, steep slopes, and proximity to water (particularly in the  
15 southern Sierra Nevada) (Zielinski et al. 2004b, Carroll 2005).

16  
17 The relatively strict reliance of fisher on dense canopy forest and complex vegetation structure,  
18 and its preference for biologically productive landscapes at lower to middle elevations has led to  
19 increasing conflicts with human habitat use and management (Powell 1993, Harrison et al. 2004,  
20 Zielinski et al. 2005). Although much research has been carried out on fisher biology, fisher  
21 habitat, and fisher distribution, no assessment of the potential direct or indirect impacts of future  
22 climate change on the California fisher has yet been published. For example, climate-driven  
23 changes in tree species dominance at fisher sites may decrease - or increase - the amount of  
24 habitat available for foraging, denning or resting. Direct impacts of climate change on fisher  
25 metabolism and reproduction are also likely. The potential indirect impacts of climate-change  
26 through climate-driven changes in fire regimes are also important to evaluate. Fire regimes  
27 respond rapidly to changes in climate and – along with other ecological disturbances like insects  
28 and disease outbreaks – are likely to drive much of the short-term response in vegetation floristics  
29 and structure (Flannigan et al. 2000, Dale et al. 2001). If longer or more severe fire seasons are  
30 one outcome of climate warming, the probability of losing local populations of species that  
31 depend on late seral habitat will increase (McKenzie et al. 2004). This is particularly a concern  
32 for species like the fisher which have relatively small range sizes and low abilities or tendencies  
33 to disperse.

#### 34 35 *Future climate change in California*

36  
37 In this contribution, results from recent modeling efforts and scientific publications are used to  
38 provide some guidance as to the likely effects of future climate change on California fisher  
39 habitat. The intent of this treatment is not to focus on direct effects of climate on fisher (e.g., on  
40 metabolic rates), but rather to make an estimation of impacts we might expect to see on  
41 vegetation composition, fire frequency and intensity, forest structure, etc., all of which have  
42 important indirect effects on fisher survival. This contribution will not provide a general  
43 overview of climate change predictions for California or the West Coast: dozens of climate  
44 models and papers exploring their implications have been published, and it is not the intent nor  
45 the place of this paper to summarize these efforts for the reader (for general overviews, see, e.g.,  
46 Field et al. 1999, NAST 2000, IPCC 2001, EPRI 2003, Wilson et al. 2003). In general, it can be  
47 said that California appears to be especially vulnerable to climate change impacts. This is due,  
48 among other things, to California's geographic location and longitudinal alignment along the  
49 Pacific Coast; its Mediterranean climate, which drives extreme contrasts in summer and winter  
50 precipitation patterns; its extremely high diversity of ecosystems and species; its large and rapidly

1 growing human population; and its international importance as an economic and agricultural  
2 powerhouse (Field et al. 1999, Snyder et al. 2002, EPRI 2003, Wilson et al. 2003).

3  
4 To this point, very few future-climate modeling efforts have treated areas as small as the State of  
5 California. The principal limiting factor is the spatial scale of the General Circulation Models  
6 (GCMs) that are used to game future climate scenarios. Most GCMs produce raster-based  
7 outputs with pixels that are 10,000's of km<sup>2</sup> in area; for example, the Hadley model referenced  
8 below produces output on a 2.5° by 3.75° (latitude by longitude) grid. To be used at finer scales,  
9 these outputs must be “downscaled” using a series of algorithms and assumptions – these finer-  
10 scale secondary products currently provide the most credible sources we have for estimating  
11 potential outcomes of long-term climate change for California. Another complication is the  
12 extent to which GCMs disagree with respect to the likely outcomes of climate change. For  
13 example, a recent comparison of 21 published GCM outputs that included the California region  
14 found that estimates of future precipitation ranged from a 26% increase per 1° C increase in  
15 temperature to an 8% decrease (Gutowski et al. 2000, Hakkarinen and Smith 2003). That said,  
16 there was some broad consensus: all of the reviewed GCMs predicted warming temperatures, and  
17 13 of 21 predicted higher precipitation (three showed no change and five predicted decreases).  
18 Most models also agreed that summers will be drier than they are currently, regardless of levels  
19 of annual precipitation.

20  
21 Currently the only modeling effort which has directly linked GCM outputs with subroutines  
22 modeling impacts on terrestrial vegetation and ecosystem processes across California is the effort  
23 conducted by the Electrical Power Research Institute (“EPRI”) under commission to the  
24 California Energy Commission (EPRI 2003). The EPRI project is the most detailed and  
25 comprehensive study yet undertaken on the potential effects of climate change in California.  
26 EPRI coordinated a diverse group of researchers and modelers who carried out a series of studies  
27 focusing on climate change impacts on terrestrial vegetation, biodiversity, timber production,  
28 water resources, agriculture, energy and coastal resources. The EPRI project also considered a  
29 number of climate change scenarios, ranging from much warmer and wetter than today, to  
30 somewhat warmer and drier than today. The project was designed to help California natural  
31 resource managers and other policy makers better understand the potential effects of climate  
32 change on the state, so as to allow the development of adaptive land and resource management  
33 policies (EPRI 2003).

34  
35 Based on the uncertain outcomes of climate change for California, one of the principal purposes  
36 of the EPRI project was to game a number of future scenarios using different estimates of change  
37 in temperature and precipitation. The EPRI team (EPRI 2003) found that the Hadley CM2  
38 scenario (Johns et al. 1997; this model was also used in the U.S. National Assessment [NAST  
39 2000]) produced one of the wettest future scenarios, and it was also the most accurate of the  
40 reviewed “wet future” models with respect to its ability to reproduce current precipitation patterns  
41 (Hakkarinen and Smith 2003). The Hadley CM2 GCM (hereafter “HAD”) was therefore chosen  
42 to represent the wet end of the modeling spectrum. The National Center for Atmospheric  
43 Research parallel climate model (hereafter “PCM”; Dai et al. 2001) produced one of the driest  
44 future scenarios, and was chosen to represent the dry end of the spectrum; a number of  
45 intermediate scenarios were also generated to represent the middle ground of GCM predictions  
46 for California. Both the HAD and PCM models are “state-of-the-art” GCMs, and include the  
47 influences of ocean dynamics and aerosol forcing on the atmosphere (EPRI 2003). For  
48 calibration, both GCM models were run from the 1800s to 1995 using observed increases in  
49 greenhouse gas concentrations. Future scenarios were based on Intergovernmental Panel on  
50 Climate Change (IPCC) projections of a 1% increase in greenhouse gases per year (EPRI 2003).

1 Finally, the EPRI project used a standard downscaling methodology to create higher resolution  
2 (100 km<sup>2</sup>) outputs from the coarse-scale GCM scenarios.

3  
4 For the purposes of this study, the most germane results from the EPRI project are those  
5 pertaining to terrestrial ecosystems (Lenihan et al. 2003a,b). Lenihan et al. used a dynamic  
6 ecosystem model (“MC1”) which estimates the distribution and the productivity of terrestrial  
7 ecosystems such as forests, grasslands, and deserts across a grid of 100 km<sup>2</sup> cells. To this date,  
8 this is the highest resolution at which a model of this kind has been applied in California. Based  
9 on their modeling results, Lenihan et al. (2003a,b) projected that, under all modeled climate  
10 change scenarios, forest types and other vegetation dominated by woody plants in California  
11 would migrate to higher elevations as warmer temperatures make those areas suitable for  
12 colonization and survival. For example, with higher temperatures and a longer growing season,  
13 the area occupied by subalpine forests and alpine vegetation was predicted to decrease as  
14 evergreen conifer forests and shrublands migrate to higher altitudes. Under their “wet future”  
15 scenarios, Lenihan et al. projected a general expansion of forests in northern California and  
16 grasslands in southern California. With higher rainfall and higher nighttime minimum  
17 temperatures, broadleaf trees were predicted to expand their distribution in many parts of the  
18 Sierra Nevada and Klamath Ranges, and conifer-dominated forests were predicted to decrease in  
19 extent in the same areas. Under their “dry future” scenarios, Lenihan et al. (2003a,b) predicted  
20 that grasslands would expand throughout the state, and that increases in the extent of tree-  
21 dominated vegetation would be minimal. Lenihan et al. also found that the frequency and the size  
22 of fires would increase under most of their climate scenarios; the change became most  
23 pronounced toward the end of the century. The drier climate scenarios resulted in more frequent  
24 fires and more area consumed by fires. Modeling based on the wetter climate scenarios suggested  
25 that fires under these conditions would tend occur at longer return intervals but at higher  
26 intensities due to greater productivity and more rapid growth of fuels.

27  
28 Studies of fisher ecology have generally found that fisher survival is less tied to forest floristics  
29 than to forest structure (e.g., Powell 1993; Zielinski et al. 2004a,b, 2006). Due to the geographic  
30 scale of their effort, Lenihan et al. (2003a,b) were not able to assess likely changes in vegetation  
31 structure in their modeling effort. I used simulation modeling of forest growth and fire to assess  
32 the possible outcomes of high, mixed and low severity fire on the kinds of forest stands preferred  
33 by fisher in the southern Sierra Nevada. In this case, my interest was in quantifying possible  
34 changes in measures of forest structure that correlate closely with fisher presence and/or density  
35 (Seglund 1995, Klug 1997, Carroll et al. 1999; Zielinski et al. 2004a,b; Carroll 2005, Yaeger  
36 2005). In this contribution, I summarize the results of the EPRI (2003) modeling efforts for the  
37 two areas currently inhabited by California fisher, present the results of vegetation growth and  
38 disturbance simulations, and discuss the possible outcomes of predicted climate change for the  
39 fisher and its habitat.

## 40 41 METHODS

### 42 43 *Summary of climate and terrestrial vegetation modeling outputs*

44  
45 Mapped outputs from the EPRI (2003) and Lenihan et al. (2003a,b) modeling efforts were  
46 imported into a graphics program. The following outputs were accessed: temperature and  
47 precipitation; vegetation type (see Table 1), fire return interval (FRI, in years and % change over  
48 time), and fireline intensity (in btu/ft/sec and % change over time). Polygons representing the  
49 approximate extent of the two California populations of fisher (drawn based on information in  
50 Zielinski et al. 2004a and 2005b) were overlain onto the EPRI and Lenihan et al. maps (Figure 1).  
51 In the graphics program, a grid was created to match the grid size in the map outputs (10 km by

1 10 km). Using the grid, map pixels within the boundaries of each fisher population were counted  
 2 and tallied as to attribute. Pixels which fell on the polygon edge were counted as  $\frac{3}{4}$ ,  $\frac{1}{2}$ , or  $\frac{1}{4}$ ,  
 3 depending on approximate area within the polygon. The number of pixels of each mapped  
 4 attribute was divided by the total number of pixels and multiplied by 100 to give percent-of-  
 5 mapped-area. Outputs for each fisher population area were graphed, using the PCM and HAD  
 6 climate model results (see above) in order to bracket the range of reasonably probable future  
 7 climate scenarios.

### 8 9 *Fire effects simulations*

10  
11 The Fire and Fuels Extension to the Forest Vegetation Simulator (FFE-FVS) (Reinhart and  
 12 Crookston 2003) was used to model fire effects on the structure of late-seral conifer vegetation in  
 13 the southern Sierra Nevada. Base data were the set of FIA plots sampled on the Sierra and  
 14 Sequoia National Forests, stratified to represent two vegetation types: mixed conifer stands  
 15 (multiple conifer species with black oak) with cover  $\geq 40\%$  and overstory trees averaging 24-39  
 16 inches (61-99 cm) diameter-at-breast-height; and fir-dominated stands with cover between 40 and  
 17 69% and overstory trees averaging 24-39 inches. California Forest Service codes for vegetation  
 18 classes making up the two types are: "M4N" and "M4G" for the mixed conifer type (I will refer  
 19 to this type as "M4N/G"), and "F4N" for the fir type. The sample size for each vegetation type  
 20 was 9.

21  
22 Computer simulations were used to grow trees in each plot for 40 years, beginning in 2006,  
 23 running simulations for high (13 ft [4 m] flame length), mixed (6.5 ft [2 m]) and low (3.3 ft [1  
 24 m]) severity fire in the first year (i.e., fires were simulated only in 2006). As a control, the same  
 25 stands were also simulated without fire. The following outputs were tracked: tree canopy cover;  
 26 number of "large trees" (dbh  $\geq 30$  inches [76 cm]) per acre; number of hardwood trees (dbh  $\geq 11$   
 27 inches [28 cm]) per acre; number of large snags (dbh  $\geq 30$  inches [76 cm]) per acre; and down  
 28 logs  $\geq 12$  inch (30 cm) diameter, measured as tons per acre. Each of these variables has been  
 29 shown to correlate with occupied fisher habitat in California and elsewhere (Powell 1993,  
 30 Zielinski et al. 2004a,b, 2006). FFE-FVS pools plots within each class before simulating growth  
 31 and disturbance, which results in a single output prediction and makes statistical comparisons  
 32 among plot outcomes difficult. Given the broad, informational purposes of this contribution, I  
 33 will rely on simple description of the simulation differences.

## 34 35 RESULTS

### 36 37 *Review of EPRI (2003) climate and ecosystem modeling results for California fisher population* 38 *areas*

#### 39 40 Climate

41 Temperature and precipitation predictions for both climate change scenarios for the period 2000-  
 42 2099 are presented in Figure 2. On the scale of the entire state, the HAD model predicts a 3.3° C  
 43 average rise in temperature by 2099, while the PCM model expects a 2.4° C rise. Averaged  
 44 across the state, the HAD model predicts an increase of about 58% in mean annual precipitation,  
 45 the PCM model a 21% decrease (Figure 2; EPRI 2003). Within the fisher population areas, the  
 46 HAD model predicts increases of about 3.1 to 3.2° C (with slightly more warming in the southern  
 47 Sierra Nevada), and the PCM model predicts increases of 2 to 2.2° C. Predicted values for mean  
 48 annual precipitation are very different between the two GCMs: by 2099, the HAD model expects  
 49 a 30-50% increase in precipitation in the general Klamath region, and a 50-70% increase in the  
 50 southern Sierra Nevada (with increases rising to the south). The PCM model predicts decreases

1 of 10-15% in the Klamaths (with some small patches of 16-20% decreases), and 5 to 15% in the  
 2 southern Sierra Nevada (EPRI 2003).

3  
 4 The EPRI (2003) outputs allow the differentiation of predicted changes in precipitation into  
 5 seasonal components (Figure 3). Within the area of the two California fisher populations, the  
 6 HAD future climate scenario (warmer and wetter) expects significant increases in wet-season  
 7 (November-March) precipitation ranging from 150 to 200% by the period 2080-2099. Within  
 8 (most of) the Klamath area, dry season (June-August) precipitation during this period is predicted  
 9 by the HAD model to be between 75 and 100% of current, but dry season precipitation is  
 10 expected to increase in the southern Sierra Nevada due to increased thunderstorm activity (Figure  
 11 3; EPRI 2003). The PCM scenario (warmer and drier) for the Klamath area predicts slightly  
 12 decreased precipitation for the wet season, but only 50-75% of current precipitation during the  
 13 dry season. The PCM predictions for the southern Sierra are similar, but expect less relative loss  
 14 in summer precipitation than in the Klamaths (Figure 3; EPRI 2003).

### 15 16 Vegetation

17 Figure 4 gives modeling results from Lenihan et al. (2003a,b) for vegetation types within the two  
 18 fisher population areas. In the Klamath area, current vegetation is modeled by MC1 as about  
 19 78% evergreen conifer forest, which includes a wide variety of conifer-dominated vegetation  
 20 types (Table 1), and about 16% mixed evergreen forest (oaks and other hardwoods in  
 21 combination with Douglas-fir or Ponderosa Pine); the remaining 5% is modeled as shrublands or  
 22 “mixed evergreen woodland” (mostly xeromorphic oak-dominated vegetation, such as blue oak  
 23 savanna or canyon live oak stands). The HAD future climate scenario (warm and wet) predicts a  
 24 reversal in the dominance of conifer- versus hardwood-dominated vegetation types in this part of  
 25 California: evergreen conifer forest suffers a +/- 90% loss in area and mixed evergreen forest  
 26 increases by almost 450%. The PCM future climate scenario (warm and dry) also predicts losses  
 27 in evergreen conifer forest (70% loss in area) and gains in hardwood-dominated vegetation  
 28 (250% increase), with significant gains in shrublands and mixed evergreen woodland as well  
 29 (Figure 4A).

30  
 31 In the southern Sierra Nevada, MC1 models about 25% of the current landscape as evergreen  
 32 conifer forest, about 22% as mixed evergreen woodland, 18% as subalpine/alpine, 17% as  
 33 shrubland, 16% as grassland, and 2% as mixed evergreen forest (Figure 4B; Lenihan et al.  
 34 2003a,b). Under the HAD future climate scenario, the area occupied by mixed evergreen forest  
 35 rises dramatically, to more than 60% of the landscape, replacing evergreen conifer forest at higher  
 36 elevations (which drops by 50% in area) and mixed evergreen woodland at lower elevations  
 37 (which loses 42% of its current area); all of the other vegetation types also drop in area. Under  
 38 the PCM scenario, the amount of mixed evergreen forest also rises, but only to about 10% of the  
 39 area; the big winner is grassland, which expands dramatically due to higher temperatures, greater  
 40 drought and higher fire frequencies (Figure 4B; Lenihan et al. 2003a,b).

### 41 42 Fire return intervals (FRI)

43 Under the HAD climate change scenario, the MC1 model (Lenihan et al. 2003a,b) predicts that  
 44 about 40% of the Klamath landscape will experience fire less often than under “current” (mean of  
 45 1895-1994) conditions, with most of this landscape seeing increases in FRI of 10-25%; under this  
 46 scenario about 50% of the landscape will see no significant change in FRI (Figure 5A). For the  
 47 same area, the drier PCM scenario results in more frequent fire over about half of the landscape,  
 48 with most of these areas seeing decreases of about 10-25% in FRIs (Figure 5A; Lenihan et al.  
 49 2003a,b). The HAD and PCM climate change scenarios make much more congruent predictions  
 50 for the southern Sierra Nevada: in both cases MC1 expects around 65% of the landscape to  
 51 experience similar FRIs to “current” conditions, and in both cases most of the remaining

1 landscape is predicted to see more frequent fire; under the HAD scenario about 11% of the  
2 southern Sierra Nevada area sees less fire (Figure 5B).

#### 3 4 *Fireline intensities*

5 As with FRIs, the HAD and PCM scenarios within the MC1 model generated disparate outcomes  
6 for the Klamath area but more or less similar outcomes for the southern Sierra Nevada (Figure 6).  
7 In the Klamaths, MC1 predicts that – under the HAD climate scenario – about ¼ of the area will  
8 see increased fire intensities by 2099 (due primarily to more fuels growth under wetter  
9 conditions), about 60% of the area will see more or less no change, and about 10% will see  
10 decreased fire intensities; the PCM-based outputs for the same area predict that about 50% of the  
11 area will see decreased fire intensities due to more frequent fire and vegetation type shifts to more  
12 flammable fuels (Figure 6A). In the southern Sierra Nevada, the HAD scenario results in about  
13 40% of the area seeing increased fire intensities, 21% experiences decreased intensities, and 38%  
14 experiences no measurable change. Landscape patterns under the PCM scenario are very similar:  
15 35% of the southern Sierra Nevada area is predicted to have fires of greater intensities by 2099,  
16 13% of the area has lower intensities, and about ½ the area is unchanged (Figure 6B; Lenihan et  
17 al. 2003a,b).

#### 18 19 *Growth and Fire Effects Simulations*

20  
21 Figures 7 and 8 report quantitative results from the FFE-FVS simulations for the mixed conifer  
22 type (classes M4N/G); results for the fir type (F4N) are not shown, as they were very similar. It  
23 is important to remember that the results are dependent on the original set of plots entered into the  
24 simulation – the reader should focus on the relative changes between years and “treatments”  
25 rather than on the actual values. Figure 7 reports results for large trees, canopy cover, and  
26 hardwood trees. The number of large trees ( $\geq 30$  inch dbh) was reduced to nearly zero by the high  
27 severity burn, mixed severity fire resulted in a 50% loss of large trees, while low severity caused  
28 a minor loss in large tree density (Figure 7A). Based on the modeled growth rates, it will be  
29 many decades beyond 2046 before there is any recovery in large tree density in the high severity  
30 simulation. In 40 years, overstory cover increased from 61% to 78% in the undisturbed  
31 simulation; rates of cover increase were much more rapid in the burned simulations. Even with  
32 an 82% loss in canopy after fire, the high severity simulation matched the undisturbed simulation  
33 in cover after 30 years; the low and mixed severity simulations were not far behind (Figure 7B).  
34 Canopy cover in the high severity simulation was contributed almost entirely by small trees.  
35 Between 2026 and 2046, ratios between tree size classes went from 56:1:0.2, to 1:1:0.2, to 1:2:0.2  
36 (6-11 inch dbh: 12-24 dbh:  $>24$ ” dbh); smaller trees contributed most cover to the mixed severity  
37 simulation as well (data not shown). The results for hardwoods (Figure 7C) are difficult to  
38 extrapolate and of limited usefulness, as the mixed conifer plots used in the simulation had no  
39 hardwoods greater than 11 inch dbh before the simulated disturbances. As a result, both high and  
40 mixed severity fire removed all hardwood trees from the simulated plots and recruitment and  
41 growth was not rapid enough to replace any of this loss before 2046 (Figure 7C); even low  
42 severity fire reduced the hardwood density by 2/3. Had there been larger hardwood trees in the  
43 plots, low and mixed severity fire would have reduced conifer cover and likely resulted in a  
44 positive hardwood growth response. Hardwood numbers in the undisturbed simulation actually  
45 dropped over time, due to increased density of conifers; again, this is also partly the result of the  
46 small sizes of hardwoods in the M4N/G plots.

47  
48 Figure 8 reports simulation results for large snags ( $\geq 30$  inch dbh) per acre and down logs ( $\geq 12$ ”  
49 dbh), measured in tons per acre. Burning of all kinds increased snags and down logs over the  
50 undisturbed simulation. Based on background rates of large tree mortality, the undisturbed  
51 simulation predicted between 4 and 5 large snags per acre by 2046. Snag creation was greatly

1 enhanced by fire, especially high intensity fire, but the total snag numbers began to converge  
 2 across simulations as time passed and snags fell. Even so, 40 years after fire the high severity  
 3 simulation still had 75% more snags than the undisturbed simulation (Figure 8A). As snags fall,  
 4 they become down woody debris. Figure 7B shows that falling trees after fire significantly  
 5 increased the amount of woody debris through the period of the simulation. The different  
 6 simulations were approximately equal immediately after fire, but had diverged significantly by  
 7 2016. The rate of woody debris creation was dependent on the severity of fire (Figure 8B), and  
 8 peaked about 20 years after the event, when there was about 14 times as much down woody  
 9 debris in the high severity simulation as in the unburned simulation (14.6 tons vs. 1.2 tons per  
 10 acre).

11  
 12 Figure 9 shows stand-level results of the FFE-FVS simulations for the mixed conifer type  
 13 (M4N/G classes) as pictured by the Stand Visualization System (McGaughey 1997). The first  
 14 column in Figure 9 is from 2006, the beginning year of the simulations; the second column is  
 15 from 2026, halfway through the simulation. Remember that fires were only burned in 2006 – the  
 16 2026 stands are thus always the results of 20 years of undisturbed growth. The figure is  
 17 organized with the average undisturbed plot in the top row, the average plot burned at high  
 18 severity in the second row, the average plot burned at mixed severity fire in the third, and the  
 19 average plot burned with low severity fire in the bottom row (Figure 9). Note the loss of canopy  
 20 cover in the high severity burn, the loss of hardwoods in the high and mixed severity burns, and  
 21 the high rate of snag and down log generation in the high and mixed severity burns as well  
 22 (Figure 9).

## 23 24 DISCUSSION

25  
 26 It is impossible to say with certainty what future climates will look like in California. That said,  
 27 there is nearly unanimous agreement that atmospheric CO<sub>2</sub> levels will more than double and  
 28 temperatures will rise by at least a few degrees C by the end of the 21<sup>st</sup> century (Field et al. 1999,  
 29 IPCC 2001). Most regional modeling efforts suggest that mean annual precipitation across  
 30 California will remain similar to current values or increase (with strong regional differences),  
 31 rain:snow ratios will increase, and the seasonality of precipitation in the state will become more  
 32 pronounced (Gutowski et al. 2000, EPRI 2003, Hakkarinen and Smith 2003, Hayhoe et al. 2004).  
 33 It is also impossible to say with certainty how climate change will impact the California fisher or  
 34 its habitat. This is due to a number of factors, chief among them: (1) differing GCM predictions  
 35 with respect to future precipitation patterns in California; (2) the broad spatial scale of current  
 36 GCM and derivative models, which cannot properly model the relatively fine-grained topographic  
 37 and forest structural landscapes that support fisher populations in California; and (3) a lack of  
 38 information as to the likely direct impacts of climate change on fisher metabolism. As with  
 39 climate, however, it appears we can make a few useful generalizations about the probable future  
 40 state of fisher habitat in California.

41  
 42 Results of the EPRI modeling effort (EPRI 2003; Lenihan et al. 2003a,b) suggest that there may  
 43 be important qualitative similarities in vegetation response to climate change in those parts of the  
 44 Klamath Mountains and southern Sierra Nevada currently supporting fisher populations. For  
 45 both areas, under both the HAD and PCM climate change scenarios, warming temperatures and  
 46 increasing CO<sub>2</sub> are predicted to result in increased growth rates and – with the distribution of  
 47 proportionally more precipitation in the winter (Miller et al. 2003a,b) – increased overall  
 48 dominance of deeply-rooted woody lifeforms, especially trees (Lenihan et al. 2003a,b). In both  
 49 areas, Lenihan et al. (2003a,b) also expect mixed evergreen forest to partially replace evergreen  
 50 conifer forest, especially where precipitation increases are large. This is due to the positive  
 51 impacts of warming and increased water availability on survival and competitiveness of broadleaf



1 (especially deciduous) hardwoods vis-à-vis evergreen conifers (Woodward 1987, Lenihan et al.  
 2 2003a,b), which will probably be forced to migrate to higher elevations to track environmental  
 3 conditions (Shafer et al. 2001). Under the wetter HAD scenario, mixed evergreen forest is also  
 4 predicted to expand to lower elevations into area currently occupied by shrubland and mixed  
 5 evergreen woodland (in this case, primarily canyon live oak, which tends to dominate south-  
 6 facing canyon slopes under current climates). Hayhoe et al. (2004), basing their analysis on more  
 7 detailed information on greenhouse gas emissions, also predicted a general statewide pattern of  
 8 largescale shifts from evergreen conifer forest to mixed evergreen types. In the southern Sierra  
 9 Nevada, the principal difference between the GCM scenarios in Lenihan et al. (2003a,b) is the  
 10 high proportion of future landscape in grassland under the drier PCM model. This transition is  
 11 caused by a decline in dormant season precipitation and an increase in frequency of fire, but  
 12 primarily affects shrublands and areas of mixed evergreen woodland below the core range of  
 13 fisher. Hayhoe et al. (2004) reported similar results under their dry future scenario.

14  
 15 Lenihan et al.'s (2003a,b) results can be compared to those of Miller and Urban (1999), who used  
 16 a different model to predict forest and fire response to climate change in the southern Sierra  
 17 Nevada. Like the EPRI effort, Miller and Urban (1999) examined responses to a variety of  
 18 different climate change scenarios, but along an elevation gradient at the geographic coordinates  
 19 of Sequoia National Park. Only the two lowest sites modeled (1800 and 2200 m) are found  
 20 within elevations currently frequented by fisher. Miller and Urban's results are also from the  
 21 period 500 to 800 years in the future, and are thus not directly comparable with the 100 year  
 22 outputs from the EPRI project. Under all climate scenarios (excepting their highly unlikely  
 23 "cooler and wetter" scenario), woody biomass decreased at both "low elevation" sites: at 1800 m,  
 24 biomass ultimately decreased to nearly zero, and at 2200 m it decreased by an average of about  
 25 70% (Miller and Urban 1999). This biomass loss was accompanied by a shift in tree species  
 26 composition at the two sites: at 1800 m, the modeled vegetation shifted from forest to grassland  
 27 and shrubland, with a small remnant component of ponderosa pine, black oak and/or incense  
 28 cedar, depending on the climate scenario. At 2200 m, species dominance shifted from white fir to  
 29 incense cedar and ponderosa pine. Miller and Urban (1999) also found that fires were more  
 30 frequent under future climate at all modeled elevations.

31  
 32 A number of previous studies have underlined the importance of hardwoods (especially large  
 33 hardwoods) to California fisher resting and denning habitat (see, e.g., Carroll et al. 1999,  
 34 Zielinski et al. 2004b). A study commissioned by the US Forest Service in California found that  
 35 fisher presence in the Klamath region began to decrease when the conifer proportion of the tree  
 36 canopy began to exceed 55% (Carroll 2005). The increase in mixed evergreen forest predicted  
 37 for both the Klamath and southern Sierra Nevada by Lenihan et al. (2003a,b) suggests that  
 38 *floristic* conditions for fisher survival may actually be enhanced by climate warming, as long as  
 39 annual precipitation remains near or above current levels, and as long as direct temperature (or  
 40 other) effects on the metabolisms of fisher and its prey do not force significant uphill movement.  
 41 For example, stands currently dominated by white fir-ponderosa pine forest might transition in  
 42 part to ponderosa pine-black oak, while Douglas fir-white fir forests (primarily in the Klamaths)  
 43 might be partially replaced by Douglas fir-tanoak (Lenihan 2003a,b). In the case of Oregon white  
 44 oak (*Quercus garryana*), a decrease in suitable habitat in the western Klamaths may be offset by  
 45 enhanced climatic suitability in the Sierra Nevada (Shafer et al. 2001). Many researchers are  
 46 expecting a future tendency toward more frequent fire throughout montane California (see below;  
 47 Field et al. 1999, Miller and Urban 1999, Lenihan et al. 2003a,b). This may further benefit oaks  
 48 (and other hardwoods) at the expense of conifers, as most oak tree species resprout after fire and  
 49 require periodic disturbances to set back conifer competition (Abrams 1992, Chang 1996).

50

1 Although oak species and other hardwoods might benefit from warming temperatures and  
2 increased frequencies of fire in montane California, the impacts of future climate change on  
3 conifer species and overall forest structure may be less positive. There is abundant evidence that  
4 fisher habitat use is more strongly driven by forest physical structure than by tree species  
5 composition (Powell 1993, Zielinski et al. 2004a,b, 2006), but the fact remains that almost all  
6 fisher habitat in California is conifer-dominated. This is simply because the key structural  
7 features of fisher habitat (high tree canopy cover, high densities of snags and down logs, the  
8 presence of large cavity-forming trees, etc.) include elements that are best represented in late-  
9 seral forests, and late-seral forests in California are overwhelmingly dominated by conifer  
10 species. Note that it is not at all impossible for younger forests to provide the habitat attributes  
11 necessary for fisher occupancy (Powell 1993, Lewis and Stinson 1998). For example, many of  
12 the stands in which Zielinski et al. (2004a,b) found California fisher were mid-seral, but this is  
13 probably at least partly a result of the general lack of late-seral forests in the areas they sampled.  
14

15 The combination of warmer climate with higher CO<sub>2</sub> fertilization will likely cause more frequent  
16 and more extensive fires throughout western North America (Price and Rind 1994, Flannigan et  
17 al. 2000); fire responds rapidly to changes in climate and will likely overshadow the direct effects  
18 of climate change on tree species distributions and migrations (Flannigan et al. 2000, Dale et al.  
19 2001). A temporal pattern of climate-driven increases in fire activity is already apparent in the  
20 western United States (Westerling et al. 2006), and modeling studies specific to California expect  
21 increased fire activity to persist and possibly accelerate under most future climate scenarios, due  
22 to increased growth of fuels under higher CO<sub>2</sub> (and in some cases precipitation), decreased fuel  
23 moistures from warmer dry season temperatures, and possibly increased thundercell activity  
24 (Price and Rind 1994, Miller and Urban 1999, Lenihan et al. 2003a,b; Westerling and Bryant  
25 2006). Increased frequencies and/or intensities of fire in coniferous forest in California will  
26 almost certainly drive changes in tree species compositions (Lenihan et al. 2003a,b), and will  
27 likely reduce the size and extent of late-successional refugia (USFS and BLM 1994, McKenzie et  
28 al. 2004). Thus, if fire becomes more active under future climates, there may be significant  
29 repercussions for the types of habitat favored by fisher in California.  
30

31 A key question is to what extent future fire regimes in montane California will be characterized  
32 by either more or less severe fire than is currently (or was historically) the case. Fire regimes are  
33 driven principally by the effects of weather/climate and fuel type and availability (Bond and van  
34 Wilgen 1996). 60 to 70 years of effective fire suppression in the American West have led to fuel-  
35 rich conditions that are conducive to intense forest fires that remove significant amounts of  
36 biomass (McKelvey et al. 1996, Arno and Fiedler 2005), and most future climate modeling  
37 predicts climatic conditions that will likely exacerbate these conditions. Basing their analysis on  
38 two GCMs under the conditions of doubled atmospheric CO<sub>2</sub> and increased annual precipitation,  
39 Flannigan et al. (2000) predicted that mean fire severity in California (measured by mean  
40 seasonal difficulty of control) would increase by about 10% averaged across the state. Vegetation  
41 growth models that incorporate rising atmospheric CO<sub>2</sub> show an expansion of woody vegetation  
42 on many western landscapes (e.g., Lenihan et al. 2003a,b), which could feedback into increased  
43 fuel biomass and connectivity and more intense (and thus more severe) fires. Use of  
44 paleoecological analogies also suggests that parts of the Pacific Northwest (including the Klamath  
45 Mountains) could experience more severe fire conditions under warmer, more CO<sub>2</sub>-rich climates  
46 (Whitlock et al. 2003). Fire frequency and severity (or size) are usually assumed to be inversely  
47 related (Pickett and White 1985), and a number of researchers have demonstrated this relationship  
48 for Sierra Nevada forests (e.g. Swetnam 1993, Miller and Urban 1999), but if fuels grow more  
49 rapidly *and* dry more rapidly – as is predicted under many future climate scenarios – then both  
50 severity and frequency may increase. Lenihan et al.'s (2003a,b) results for fire intensity (Figure  
51 5) predict that 1/3 or more of the southern Sierra landscape may see mean fire intensities increase

1 over current conditions by the end of the century. Up to ¼ of the Klamaths may see a similar  
2 increase if precipitation increases, but fire intensity decreases (and frequency increases) under the  
3 PCM warm-dry scenario, largely because of the replacement of evergreen forest by hardwoods,  
4 open woodlands, and shrub- and grass-dominated systems (Lenihan et al. 2003a,b), vegetation  
5 types that are not generally frequented by fisher.

6  
7 If future fires in forests occupied by fisher are indeed more intense on average, then we can  
8 expect the severity of fire effects to increase as well. Based on the FFE-FVS modeling (Figures  
9 7-9), it appears that increased frequencies of mixed and high severity fire in dense, mid- to late-  
10 seral stands of firs and mixed conifer species are likely to significantly reduce overstory cover  
11 and the density of large trees, two of the factors most closely correlated with fisher presence in  
12 California forests (Powell 1993, Zielinski 2004a,b). High severity, stand-replacing fire can  
13 render forest habitat unusable by fisher for many decades (Lewis and Stinson 1998). Decreases  
14 in overhead cover can lead to, among other things, increased drying effects from sun and wind in  
15 the dry season, and increased snow deposition in the wet season, which could seriously limit  
16 fisher mobility (Krohn et al. 1997) or force their restriction to smaller remnant patches of dense  
17 forest (where overhead canopy cover intercepts high proportions of snowfall; Weir 1995). Note  
18 that this tendency could be offset by decreasing snow-water ratios in future precipitation,  
19 depending on the elevation (Miller et al. 2003a,b). Loss of cover could also result in increased  
20 fisher losses from predation. Larger adult fisher are apparently rarely predated, but the  
21 probability of successful capture of fisher by predators such as raptors, bobcats, and mountain  
22 lions is thought to markedly increase in areas with low cover and simplified forest structure  
23 (Buck et al. 1994, Powell and Zielinski 1994, Truex et al. 1998; D. MacFarlane, pers. comm.).  
24 Decreases in large tree density could have negative effects on both denning and resting behavior  
25 of fisher, as fisher in California strongly prefer cavities in large, old trees (apparently to minimize  
26 the effects of summer heat and dryness; Zielinski 2004b). As the FFE-FVS modeling underlined,  
27 the effect of forest fire is generally to increase snags and the amount of down woody debris; fire  
28 also aids in the creation of tree cavities, especially where severities are low or patchy in effect.  
29 However, if frequencies of mixed and high severity fire increase as climates warm, there may  
30 well be a negative net effect on the amount and distribution of surface woody debris over the long  
31 term (USFWS 2006). Overall, probably the most significant outcome of potential losses in  
32 canopy cover and/or surface woody debris would be changes or reductions in the densities of  
33 fisher prey. California fishers are generalist predators (Powell 1993, Zielinski et al. 1999), hence  
34 they may be able to accommodate a certain amount of prey-base change. However, given that  
35 prey availability is a major driving factor in fisher distribution (Powell 1993, Powell and Zielinski  
36 1994), it may be that the local survival of fisher is driven more by direct and indirect effects of  
37 changing climates on other small mammals and birds than by effects on the fisher itself.

38  
39 Although I have focused on indirect effects of climate change on fisher, there will undoubtedly be  
40 significant direct effects as well. The distribution of any species is driven by multiple, interacting  
41 factors, but direct metabolic relationships with the physical environment usually form the  
42 foundation for a species' range (Krebs 1994, Brown and Lomolino 1998). It seems highly likely  
43 that the California fisher, which is fur-bearing and primarily restricted to cooler microhabitats  
44 (north-facing slopes, canyons) within its range, will be forced to move to cooler climates as  
45 temperatures rise. Simple use of lapse rates (which average 3.6°/1000 feet [6.5°/1000 m];  
46 Luetgens and Tarbuck 1982) suggests that "optimal" temperatures for fisher habitat will move  
47 upslope 500-1000 feet (150-300 m) over the next century. Often temperature extremes are more  
48 important than means in driving species distributions – Hayhoe et al. (2004) found that the  
49 probability of extreme heat events in California will undergo a marked increase as temperatures  
50 rise. Under the future climate scenarios they assessed, the "heatwave season" in California  
51 increased by 5 to 13 weeks in length, depending on location. In the Los Angeles Basin, about

1 100 miles (160 km) south of the southern end of the Sierra fisher population area, Hayhoe et al.  
2 (2004) estimated that there would be 22 more days of maximum temperatures greater than 90°F  
3 (32°C) by the year 2100. There will certainly be direct effects of increasing temperature on fisher  
4 prey as well, but these will operate in different ways on different species and the aggregate  
5 outcome on the fisher prey base is difficult to predict. There will also likely be temperature  
6 effects on fisher habitat use *within* its range, wherever it may be. For example, Weir et al. (2004)  
7 found that air temperature was a key factor in determining what resting structure was chosen by  
8 fisher: coarse woody debris was most often used under very cold temperatures, while “witches  
9 brooms” and branches were chosen more often under warmer conditions. There seem certain to  
10 be various, complicated interactions between these sorts of responses and future climates and  
11 disturbance regimes.

12  
13 Climate change is likely to precipitate a number of complicated (and largely unpredictable)  
14 outcomes. I have focused in this contribution on more immediate effects of climate change, but it  
15 may be that these are swamped or negated by unexpected interactions. Currently, much research  
16 is being done on the synergies of natural disturbance in western North America. For example, the  
17 complex relationships between rising temperatures and drought, water stress, insect and disease  
18 occurrence, and fire appear already to be driving rapid ecosystem response to climate change in  
19 western forests (Dale et al. 2001, Breashears et al. 2004). Such feedback effects among multiple  
20 disturbances are thought to be behind recent large-scale shifts in geographic distribution of  
21 ponderosa pine and piñon pine in the Southwest and California (Allen and Breashears 1998; J. H.  
22 Thorne, pers. comm.). In many cases, insect- or disease-caused mortality among conifers may  
23 benefit fisher by creating resting, denning and foraging habitat, but large inputs of standing and  
24 down dead wood – such as during pine beetle epidemics – may feedback into higher hazard of  
25 stand-replacing fire as well, which could ultimately cause the loss of these same habitat structures  
26 (Shaffer and Laudenslayer 2006, USFWS 2006). The foundation for many of these synergistic  
27 “loops” appears to be water stress in trees, as it weakens tree vigor and can lead to higher  
28 incidences of fire-, insect-, disease- and pollution-caused mortality. Most GCMs predict lower  
29 snow:rain ratios, reduced winter snowpack, earlier snowmelt, and lower dry-season streamflows  
30 (IPCC 2001, Miller et al. 2003a,b; Stewart et al. 2004), not to mention higher growing season  
31 temperatures, which will drive increases in evaporative demand. Recent data show that changes  
32 in these variables are already occurring in the predicted direction (Mote 2003, Stewart et al. 2005)

33  
34 Many authors have underlined the importance of microenvironment to fisher survival (e.g.,  
35 Powell 1993, Powell and Zielinski 1994, Buck et al. 2004, Harrison et al. 2004, Weir et al. 2004).  
36 The climate modeling efforts reported on in this paper simply do not have the resolution to  
37 reliably predict impacts of climate change on local habitat variables. In the EPRI modeling effort  
38 (EPRI 2003; Lenihan et al. 2003a,b), the minimum grid size was 100 km<sup>2</sup>, much finer resolution  
39 than any other comparable effort, but still much too coarse-grained to permit realistic assessments  
40 of direct or indirect impacts climate change on local habitat. Fishers are closely connected to  
41 topographic variables (slope complexity, aspect, distance from streams, etc.) and habitat variables  
42 (tree densities, down woody debris, etc.) that will require more refined modeling techniques, such  
43 as those used by Zielinski et al. (2004a,b) and those currently being undertaken by the US Forest  
44 Service and the Conservation Biology Institute in the southern Sierra Nevada (see  
45 <http://www.consbio.org/cbi/projects/fisher>). The next obvious step is to incorporate climatic  
46 change inputs into these sorts of habitat models. Forest growth and disturbance simulations can  
47 also provide insight into likely impacts of changing fire (and other disturbance) regimes on  
48 vegetation structure, but a direct functional link to GCM outputs is still lacking; work on such a  
49 link is presently underway (R. Neilson, pers. comm.). Hopefully such work can be accomplished  
50 soon - broad extrapolations of uncertain processes to uncertain effects will do little in the end to  
51 conserve species at the edge of extinction.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

## ACKNOWLEDGEMENTS

The author benefited from discussions with Ron Neilson (USFS, Pacific Northwest Research Station), Cheryl Carrothers and Diane MacFarlane (USFS, Pacific Southwest Region), and Dave Schmidt (The Nature Conservancy-California). Andy Taylor (USFS, Mendocino National Forest) carried out the FFE-FVS modeling.

## LITERATURE CITED

- Abrams, M. D. 1992. Fire and the development of oak forests. *BioScience* 42: 346-353.
- Allen, C. D., and D. D. Breashears. 1998. Drought-induced shift of a forest-woodland ecotone: rapid landscape response to climate variation. *Proceedings of the National Academy of Sciences* 95: 14839-14842
- Arno, S. F., and C. E. Fiedler. 2005. *Mimicking nature's fire. Restoring fire-prone forests in the West*. Island Press, Washington, DC, USA.
- Bond, W. J., and B. W. van Wilgen. 1996. *Fire and plants*. Chapman and Hall, London, England.
- Breashears, D. D., N. S. Cobb, P. M. Rich, K. P. Price, C. D. Allen, R. G. Balice, W. H. Romme, J. H. Karstens, M. L. Floyd, J. Belnap, J. J. Anderson, O. B. Myers and C. W. Meyer. 2004. Regional vegetation die-off in response to global-change-type drought. *Proceedings of the National Academy of Sciences* 102: 15144-15148.
- Brown, J. H., and M. V. Lomolino. 1998. *Biogeography*. 2nd edition. Sinauer Associates, Sunderland, MA, USA.
- Buck, S. G., C. Mullis, A. S. Mossman, I. Show, and C. Coolahan. 1994. Habitat use by fishers in adjoining heavily and lightly harvested forest. Pp. 386-376, in: S. W. Buskirk, A. S. Harestad, M. G. Raphael, and R. A. Powell (eds). *Martens, sables and fishers: biology and conservation*. Cornell University Press, Ithaca, NY, USA.
- Carroll, C., W. J. Zielinski and R. F. Noss. 1999. Using presence-absence data to build and test spatial habitat models for the fisher in the Klamath Region, USA. *Conservation Biology* 13: 1344-1359.
- Carroll, C. 2005. A reanalysis of regional fisher suitability including survey data from commercial forests in the redwood region. Report prepared for USDA Forest Service, Pacific Southwest Research Station. Klamath Center for Conservation Research, Orleans, California, USA.
- Chang, C. 1996. Ecosystem responses to fire and variations in fire regimes. Pp. \_\_\_, in: *Sierra Nevada Ecosystem Project: final report to Congress. Vol. II, Assessments and scientific basis for management options*. University of California, Centers for Water and Wildland Resources, Davis, CA, USA.
- Dai, A., G. A. Meehl, W. M. Washington, T. M. L. Wigley, and J. M. Arblaster. 2001. Ensemble simulation of 21st century climate changes: Business as usual vs. CO<sub>2</sub> stabilization. *Bulletin of the American Meteorological Society* 82: 2377-2388.

- 1  
2 Dale, V. H., L. A. Joyce, S. McNulty, R. P. Neilson, M. P. Ayres, M. D. Flannigan, P. J. Hanson,  
3 L. C. Irland, A. E. Lugo, C. J. Peterson, D. Simberloff, F. J. Swanson, B. J. Stocks, and B. M.  
4 Wotton. 2001. Climate change and forest disturbances. *BioScience* 51: 723-734.  
5  
6 EPRI (Electric Power Research Institute). 2003. *Global climate change and California:  
7 potential implications for ecosystems, health, and the economy*. Report commissioned by the  
8 California Energy Commission. Palo Alto, California, USA.  
9  
10 Field, C., G. Daily, F. Davis, S. Gaines, P. Matson, J. Melack, and N. Miller. 1999. *Confronting  
11 climate change in California: ecological impacts on the Golden State*. Report of the Union of  
12 Concerned Scientists and the Ecological Society of America. UCS Publications, Cambridge,  
13 Massachusetts, USA.  
14  
15 Flannigan M. D., B. J. Stocks, and B. M. Wotton. 2000. Climate change and forest fires.  
16 *Science of the Total Environment* 262: 221–229.  
17  
18 Gutowski, W. J., Z. Pan, C. J. Anderson, R. W. Arritt, F. Otieno, E. S. Takle, J. H. Christensen,  
19 and O. B. Christensen. 2000. What RCM data are available for California impacts modeling?  
20 California Energy Commission Workshop on Climate Change Scenarios for California, 12-13  
21 June, 2000. California Energy Commission, Sacramento, CA, USA.  
22  
23 Hakkarinen, C., and J. Smith. 2003. Appendix I. Climate scenarios for a California Energy  
24 Commission study of the potential effects of climate change on California: summary of a June  
25 12-13, 2000, workshop. 38 pp., in: *Global climate change and California: potential implications  
26 for ecosystems, health, and the economy*. EPRI (Electric Power Research Institute), Palo Alto,  
27 CA, USA.  
28  
29 Harrison, D. J., A. K. Fuller, and G. Proulx. 2004. *Martens and fishers (*Martes*) in human-  
30 altered environments: an international perspective*. Springer, New York, New York, USA.  
31  
32 IPCC (International Panel on Climate Change). 2001. *Climate change 2001. Third assessment  
33 report of the Intergovernmental Panel on Climate Change*. Cambridge University Press,  
34 Cambridge, UK.  
35  
36 Johns, T. C., R. E. Carnell, J. F. Crossley, J. M. Gregory, J. F. B. Mitchell, C. A. Senior, S. F. B.  
37 Tett, and R. A. Wood. 1997. The Second Hadley Centre Coupled Ocean-Atmospheric GCM:  
38 Model description, spinup and validation. *Climate Dynamics* 13: 103-134.  
39  
40 Klug, R. R. 1997. Occurrence of Pacific fisher (*Martes pennanti*) in the Redwood Zone of  
41 northern California and the habitat attributes associated with their detections. MS thesis,  
42 Humboldt State University, Arcata, CA, USA.  
43  
44 Krebs, C. J. 1994. *Ecology. The experimental analysis of distribution and abundance*. Harper  
45 Collins, New York, NY, USA.  
  
46 Krohn, W. B., W. J. Zielinski, and R. B. Boone. 1997. Relations among fishers, snow, and  
47 martens in California: results from small-scale spatial comparisons. Pp. 211-232, in: G. Proulx,  
48 H. N. Bryant, and P. M. Woodard (eds). *Martes: taxonomy, ecology, techniques, and  
49 management*. Provincial Museum of Alberta, Edmonton, Canada.

- 1 Lenihan, J. M., R. Drapek, R. P. Neilson and D. Bachelet. 2003a. Appendix IV. Climate  
2 change effects on vegetation distribution, carbon stocks, and fire regimes in California. 60 pp.,  
3 in: *Global climate change and California: potential implications for ecosystems, health, and the*  
4 *economy*. Electric Power Research Institute, Palo Alto, California, USA.
- 5 Lenihan, J. M., R. Drapek, D. Bachelet and R. P. Neilson. 2003b. Climate change effects on  
6 vegetation distribution, carbon, and fire in California. *Ecological Applications* 13: 1667-1681.  
7
- 8 Lewis, J. C., and D. W. Stinson. 1998. *Washington State status report for the fisher*.  
9 Washington Department of Fish and Wildlife, Olympia, WA, USA.
- 10
- 11 Luetgens, F. K., and E. J. Tarbuck. 1982. *The atmosphere. An introduction to meteorology*. 2<sup>nd</sup>  
12 edition. Prentice-Hall, Englewood Cliffs, NJ, USA.
- 13
- 14 Mazzoni, A. K. 2002. Habitat use by fishers (*Martes pennanti*) in the southern Sierra Nevada.  
15 MS thesis, California State University, Fresno, CA, USA.
- 16
- 17 McGaughey, R. J. 1997. Visualizing forest and stand dynamics using the stand visualization  
18 system. *Proceedings of the 1997 ACSM/ASPRS Annual Convention and Exposition*. American  
19 Society for Photogrammetry and Remote Sensing, Bethesda, MD, USA.
- 20
- 21 McKelvey, K.S., C. N. Skinner, C. Chang, D. C. Erman, S. J. Husari, D. J. Parsons, J. W. van  
22 Wagtendonk, J.W., and C. W. Weatherspoon. 1996. An overview of fire in the Sierra Nevada.  
23 Pp. 1033-1040, in: *Sierra Nevada Ecosystem Project: final report to Congress. Vol. II,*  
24 *Assessments and scientific basis for management options*. University of California, Centers for  
25 Water and Wildland Resources, Davis, CA, USA.
- 26
- 27 McKenzie, D., Z. Gedalof, D. L. Peterson and P. Mote. 2004. Climatic change, wildfire, and  
28 conservation. *Conservation Biology* 18: 890-902.
- 29
- 30 Miller, C., and D. Urban. 1999. Forest pattern, fire and climatic change in the Sierra Nevada.  
31 *Ecosystems* 2: 76-87
- 32
- 33 Miller, N. L., K. E. Bashford and E. Strem. 2003. Appendix VIII. Climate change sensitivity  
34 study of California hydrology. 47 pp., in: *Global climate change and California: potential*  
35 *implications for ecosystems, health, and the economy*. Electric Power Research Institute, Palo  
36 Alto, CA, USA.
- 37
- 38 Miller, N. L., K. E. Bashford, and E. Strem. 2003. Potential impacts of climate change on  
39 California hydrology. *Journal of the American Water Resources Association* 39: 771-784.  
40
- 41 Mote, P. W. 2003. Trends in snow water equivalent in the Pacific Northwest and their climatic  
42 causes. *Geophysical Research Letters* 30(12): pp. 3-1, CitelID 1601, DOI 10.1029/  
43 2003GL017258.
- 44
- 45 NAST (National Assessment Synthesis Team). 2000. Climate change impact on the United  
46 States: the potential consequences of climate variability and change. National Assessment  
47 Synthesis Team. U.S. Global Change Research Program, Washington, DC, USA.
- 48
- 49 Pickett, S. T. A., and P. S. White. 1985. *The Ecology of Natural Disturbance and Patch*  
50 *Dynamics*. Academic Press, New York, NY, USA

- 1  
2 Powell, R. A. 1993. *The fisher: life history, ecology, and behavior*. 2nd edition. University of  
3 Minnesota Press, Minneapolis, Minnesota, USA.  
4
- 5 Powell, R. A., and W. J. Zielinski. 1994. Fisher. Pp. 38-73, in: L. F. Ruggiero, K. B. Aubry, S.  
6 W. Buskirk, L. J. Lyon, and W. J. Zielinski (eds). *The scientific basis for conserving forest*  
7 *carnivores: American marten, fisher, lynx, and wolverine*. General Technical Report RM-254.  
8 US Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins,  
9 Colorado, USA.  
10
- 11 Price, C., and D. Rind. 1994. The impact of a 2 x CO<sub>2</sub> climate on lightning-caused fires.  
12 *Journal of Climate* 7: 1484-1494.
- 13 Reinhardt, E., and N. L. Crookston (eds). 2003. *The Fire and Fuels Extension to the Forest*  
14 *Vegetation Simulator*. Gen. Tech. Rep. RMRS-GTR-116. USDA-Forest Service, Rocky  
15 Mountain Research Station, Ogden, UT, USA.
- 16 Seglund, A. E. 1995. The use of rest sites by the Pacific fisher. MS thesis, Humboldt State  
17 University, Arcata, CA, USA.  
18
- 19 Shafer, S. L., P. J. Bartlein, and R. S. Thompson. 2001. Potential changes in the distributions of  
20 Western North America tree and shrub taxa under future climate scenarios. *Ecosystems* 4: 200-  
21 215.  
22
- 23 Shaffer, K. E., and W. F. Laudenslayer, Jr. 2006. Fire and animal interactions. Pp. 118-144, in:  
24 N. G. Sugihara, J. W. van Wagtenonk, K. E. Shaffer, J. Flites-Kaufman, and A. E. Thode (eds).  
25 *Fire in California ecosystems*. University of California Press, Berkeley, CA, USA.  
26
- 27 Snyder, M. A., J. L. Bell, L. C. Sloan, P. B. Duffy, and B. Govindasamy. 2002. Climate  
28 responses to a doubling of atmospheric carbon dioxide for a climatically vulnerable region.  
29 *Geophysical Research Letters* 29(11): 1-4.  
30
- 31 SPSS. 2001. SPSS version 11.0. SPSS Inc., Chicago, IL, USA.  
32
- 33 Stewart, I. T., D. R. Cayan, and M. D. Dettinger. 2004. Changes in Snowmelt Runoff Timing in  
34 Western North America under a 'Business as Usual' Climate Change Scenario. *Climatic Change*  
35 62:217-232.  
36
- 37 Stewart, I. T., D. R. Cayan, and M. D. Dettinger. 2005. Changes toward earlier streamflow  
38 timing across western North America. *Journal of Climate* 18:1136-1155.  
39
- 40 Swetnam, T. W. 1993. Fire history and climate change in giant sequoia groves. *Science* 262:  
41 885-889.  
42
- 43 Truex, R. L., W. J. Zielinski, R. T. Golightly, R. L. Barrett, and S. M. Wisely. 1998. A meta-  
44 analysis of regional variation in fisher morphology, demography, and habitat ecology in  
45 California [report submitted to California Department of Fish and Game]. USDA Forest Service,  
46 Pacific Southwest Research Station, Redwood Sciences Lab, Arcata, CA, USA.  
47



- 1 USFS and BLM. 1994. *Record of Decision for amendment to Forest Service and Bureau of*  
2 *Land Management planning documents within the range of the northern spotted owl*. USDA  
3 Forest Service and USDI Bureau of Land Management, Portland, OR, USA.  
4
- 5 USFWS. 2006. Species assessment and listing priority assignment form: Fisher, West Coast  
6 distinct population segment. US Fish and Wildlife Service, Portland, OR, USA.  
7
- 8 Weir, R. D. 1995. Diet, spatial organization, and habitat relationships of fishers in south-central  
9 British Columbia. MS thesis, Simon Fraser University, Burnaby, BC, Canada.  
10
- 11 Weir, R. D., F. Corbould and A. Harestad. 2004. Effect of ambient temperature on the selection  
12 of rest structures by fishers. Pp. 187-197 in: D. J. Harrison, A. K. Fuller, and G. Proulx (eds).  
13 *Martens and fishers (Martes) in human-altered environments: an international perspective*.  
14 Springer, New York, New York, USA.  
15
- 16 Westerling, A. L., and B. Bryant. 2006. Climate change and wildfire in and around California:  
17 fire modeling and loss modeling. Report from the California Climate Change Center to the  
18 California Energy Commission. CEC-500-2006-190-SF.  
19
- 20 Westerling, A. L., H. Hidalgo, D. R. Cayan, and T. Swetnam. 2006. Warming and earlier spring  
21 increases western U.S. forest wildfire activity. *Science*: 6 July, 2006 /10.1126/ science.1128834  
22
- 23 Whitlock, C., S. L. Shafer, and J. Marlon. 2003. The role of vegetation change in shaping past  
24 and future fire regimes in the northwest U.S. and the implications for ecosystem management.  
25 *Forest Ecology and Management* 178: 5-21.  
26
- 27 Wilson, T., L. Williams, J. Smith, and R. Medelsohn. 2003. *Global climate change and*  
28 *California: potential implications for ecosystems, health, and the economy*. Publication No. 500-  
29 03-058CF, California Energy Commission, Public Interest Energy Research Environmental  
30 Area, Sacramento, CA, USA.  
31
- 32 Yaeger, J. S. 2005. Habitat at fisher resting sites in the Klamath Province of Northern California.  
33 MS thesis, Humboldt State University, Arcata, CA, USA.  
34
- 35 Zielinski, W. J., N. P. Duncan, E. C. Farmer, R. L. Truex, A. P. Clevenger, and R. G Barrett.  
36 1999. Diet of fisher (*Martes pennanti*) at the southernmost extent of their range. *Journal of*  
37 *Mammalogy* 80: 961-971.  
38
- 39 Zielinski, W. J., R. L. Truex, G. A. Schmidt, F. V. Schlexer, K. N. Schmidt, and R. H. Barrett.  
40 2004a. Home range characteristics of fishers in California. *Journal of Mammalogy* 85: 649-657.  
41
- 42 Zielinski, W. J., R. L. Truex, G. A. Schmidt, F. V. Schlexer, K. N. Schmidt, and R. H. Barrett.  
43 2004b. Resting habitat selection by fishers in California. *Journal of Wildlife Management* 68:  
44 475-492.  
45
- 46 Zielinski, W. J., R. L. Truex, F. V. Schlexer, L. A. Campbell and C. Carroll. 2005b. Historical  
47 and contemporary distributions of carnivores in forests of the Sierra Nevada, California, USA.  
48 *Journal of Biogeography* 32: 1385-1407.  
49
- 50 Zielinski, W. J., R. L. Truex, J. R. Dunk, and T. Gaman. 2006. Using forest inventory data to  
51 assess fisher resting habitat suitability in California. *Ecological Applications* 16: 1010-1025.

1  
2  
3 FIGURE CAPTIONS  
4

5 Figure 1. The approximate range of fisher in California. The two population areas are indicated.  
6

7 Figure 2. Future projected trends in mean annual temperature and mean annual precipitation,  
8 averaged across California. Projections from two future climate scenarios: the Hadley Climate  
9 Center HADCM2 model (“HAD”) and the National Center for Atmospheric Research’s Parallel  
10 Climate Model (“PCM”). Redrawn from EPRI (2003).  
11

12 Figure 3. Predicted seasonal changes in precipitation for the two fisher population areas in  
13 California, based on the mean of the period 2080-2099 compared to the mean of the period 1961-  
14 1990. Projections from the “HAD” and “PCM” future climate scenarios (see text). (A) =  
15 Klamath Mountains; (B) = southern Sierra Nevada. Data represent the percent of landscape  
16 within each population area experiencing a given change. Data from EPRI 2003.  
17

18 Figure 4. Vegetation types, current (mean of period 1961-1990) versus predicted future (mean of  
19 period 2070-2099). Projections from the “HAD” and “PCM” future climate scenarios (see text).  
20 (A) = Klamath Mountains; (B) = southern Sierra Nevada. Data represent the percent of landscape  
21 within each vegetation type. Data from Lenihan 2003a.  
22

23 Figure 5. Predicted changes in fire rotation interval (FRI) for the two fisher population areas in  
24 California, based on the mean of the period 2080-2099 compared to the mean of the period 1961-  
25 1990; negative change means that fire frequency has increased. Projections from the “HAD” and  
26 “PCM” future climate scenarios (see text). (A) = Klamath Mountains; (B) = southern Sierra  
27 Nevada. Data represent the percent of landscape within each population area experiencing a given  
28 change. Data from Lenihan 2003a.  
29

30 Figure 6. Predicted changes in fire intensity for the two fisher population areas in California,  
31 based on the mean of the period 2080-2099 compared to the mean of the period 1961-1990;  
32 negative change means that fire intensity has dropped. Projections from the “HAD” and “PCM”  
33 future climate scenarios (see text). (A) = Klamath Mountains; (B) = southern Sierra Nevada.  
34 Data represent the percent of landscape within each population area experiencing a given change.  
35 Data from Lenihan 2003a.  
36

37 Figure 7. Results of FFE-FVS forest growth and disturbance simulations for the mid- to late-seral  
38 mixed conifer vegetation type (M4N/G). (A) = number of large conifer trees ( $\geq 30$  inches dbh)  
39 per acre; (B) = canopy cover (%); (C) = numbers of hardwood trees ( $\geq 6$  inches dbh) per acre.  
40

41 Figure 8. Results of FFE-FVS forest growth and disturbance simulations for the mid- to late-seral  
42 mixed conifer vegetation type (M4N/G). (A) = numbers of large snags ( $\geq 30$  inches dbh) per acre;  
43 (B) = tons per acre of down logs  $\geq 12$  inches diameter.  
44

45 Figure 9. Results of FFE-FVS forest growth and disturbance simulations for the mid- to late-seral  
46 mixed conifer vegetation type (M4N/G). Graphic representation of stand conditions in 2006 and  
47 2026 after subjection to fires of different severities in 2006.  
48  
49  
50  
51

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

Table 1. Vegetation types simulated by the MC1 dynamic ecosystem model. Table modified from Lenihan 2003a.

Vegetation type	Description	California examples
Alpine/subalpine forest	tundra, boreal forest	alpine meadows, lodgepole pine forest, whitebark pine forest
Evergreen conifer forest	maritime temperate conifer forest, continental temperate coniferous forest	coastal redwood forest, coastal closed-cone pine forest, mixed conifer forest, ponderosa pine forest
Mixed evergreen forest	warm temperate/subtropical mixed forest	Douglas-fir–tan oak forest, tan oak–madrone–oak forest, ponderosa pine–black oak forest
Mixed evergreen woodland	temperate mixed xeromorphic woodland, temperate conifer xeromorphic woodland	blue oak woodland, canyon live oak woodland, northern juniper woodland
Grassland	C <sub>3</sub> grassland, C <sub>4</sub> grassland	valley grassland, southern coastal grassland, desert grassland
Shrubland	mediterranean shrubland, temperate arid shrubland	chamise chaparral, southern coastal scrub, sagebrush steppe
Desert	subtropical arid shrubland	creosote brush scrub, saltbrush scrub, Joshua tree woodland

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51




Figure 1

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19




Figure 2

20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51




FIGURE 3

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51




FIGURE 4



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51




FIGURE 5

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51




FIGURE 6

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51






FIGURE 7







FIGURE 8



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19



**No  
fire**





**High  
severity  
fire**



**Mixed  
severity  
fire**



**Low  
severity  
fire**



**2006**

**2026**